arXiv:1202.1610v2 [cond-mat.str-el] 26 Feb 2012

Symmetric spin-orbital-Hubbard model on honeycomb lattice: An origin of Kitaev model and zig-zag antiferromagnetic order

Yue Yu and Shaojing Qin[1]
[1]*State Key Laboratory of Theoretical physics, Institute of Theoretical Physics,*
Chinese Academy of Sciences, P.O. Box 2735, Beijing 100190, China
(Dated: February 28, 2012)

We present a single band Hubbard model on honeycomb lattice with link-symmetric spin-orbital interaction. In the large U limit, this model is mapped to a conventional antiferromangentic (AFM) Heisenberg model combining with a 'zig-zag' AFM Heisenberg one. The latter AFM order which comes from the symmetric spin-orbital coupling was favored by a recent experiment for A_2IrO_3 (A=Na, Li)[6–9]. Competition between two different AFM orders leads to a spin liquid phase characterized by Kitaev model that happens if the spin-orbital coupling has the same strength as the hopping amplitude. This deepens our understanding of the origin of the Kitaev coupling: It comes from the balance between the hopping and spin-orbital coupling but to get rid of the hopping. In a moderate U, this model has a charge-spin gapped phase with two different spinon excitations. In a weak interaction, the original gapless Dirac fermion is gapped by the spin-orbital coupling while a new gapless Dirac fermion appears at the same Dirac points.

PACS numbers: 71.20.Be, 75.25.Dk, 75.30.Et, 75.10.Jm

Kitaev model on honeycomb lattice has attracted a lot of interests because of its exact solubility, Majorana fermion excitation, non-trivial topological orders and amazing non-abelian anyon [1]. This non-abelian anyon is the key object to design a device of topological quantum computer [2]. A possible way to realize Kitaev model on optical lattice has been proposed [3]. Recently, studies of iridium oxides showed that this peculiar quantum spin liquid (SL) is possible to be realized[4, 5]. While the experiments for A_2IrO_3 (A=Na, Li) supported a 'zig-zag' antiferromagnetic(AFM) phase [6–9], Kitaev SL is possible to be observed in a high pressure [7]. These progresses drive a number of further researches [10–17] .

It is argued that the strong spin-orbtial(SO) interaction plays an important role in deriving the Kitaev-Heisenberg (KH) model [4]. However, the exact soluble Kitaev model is on the opposite side of the Neel order, i.e., one has to get rid of the hopping to obtain it.

In this Letter, we will directly show how the SO interaction induces a Kitaev coupling in the context of a single band repulsive Hubbard model. What we observe is as follows: For a large U, while the hopping between the nearest neighbor sites contributes to the Neel order of spin $\mathbf{S}$, the SO coupling offers a 'zig-zag' AFM order with a variant spin $\tilde{\mathbf{S}}$. The latter one is the most likely spin structure in A_2IrO_3 according to the recent experiments [6–9]. The competition between these two different AFM orders gives birth to a SL phase delegated by Kitaev model with anyonic excitations. In fact, when the SO coupling strength is the same as the hopping amplitude, the resulting spin model is exactly Kitaev model. This is the crucial difference of our result from that in the HK model: The Kitaev phase is in the midway between the hopping amplitude and the SO coupling but getting rid of the hopping. We notice that the Kitaev phase may also appear in the competition between two different FM phases .

The SO interaction which can lead to previously mentioned physics is a time-reversal invariant SO interaction with the nearest neighbor site, i.e., link-dependence. The SO couplings with an orientation-dependence exist in many Mott insulators with orbital orders and fluctuations[18]. In this Letter, we include a particular link-dependent SO coupling to the Hubbard model and call it the symmetric SO-Hubbard (SSO-Hubbard) model. The strong coupling Mott physics has been summarized in the previous paragraph. For a moderate U, a gapped SL phase has been predicted even for the Hubbard model without a SO coupling[19]. A slave boson theory [20, 21] also showed that there is such a SL state at the mean field level but may be broken down due to a staggered $U(1)$ gauge fluctuation for some symmetries are broken [22]. This SL is also found in a Kane-Mele-Hubbard model [23]. What we find is that, by using the slave boson method [22], the SSO coupling will enlarge the original spinon gap while a new branch of spinon excitation takes the gap of the original one. In the weak coupling limit, we also find that the well-known Dirac fermion on the honeycomb lattice is gapped by the SSO-coupling while a new gapless Dirac fermion whose velocity is solely determined by SSO coupling is exactly locates at the original Dirac points.

We consider the SSO-Hubbard model on honeycomb lattice with links $a = x, y, z$ described by the following Hamiltonian

$$\begin{aligned} H_{SSOH} &= H_T + H_{SSO} + H_U = -t\sum_{\langle ij\rangle}(c^\dagger_{i\sigma}c_{j\sigma} + h.c) \\ &+ i\lambda \sum_{a;\langle ij\rangle_a} \nu_{ij} c^\dagger_{i\sigma}\sigma^a_{\sigma\sigma'}c_{j\sigma'} + U\sum_i n_{i\uparrow}n_{i\downarrow} \qquad (1) \end{aligned}$$

where $\nu_{ij} = -\nu_{ji} = 1$. $c_{i\sigma}$ is the electron annihilation operator at site i. σ^a are Pauli matrices. $U \geq 0$ is Hubbard potential. $\langle ij \rangle$ labels the summation over the nearest neighbor sties and $\langle ij \rangle_a$ means over all the a-links. Different from that in Kane-Mele model [24], here the spin orbital coupling is x, y, z-cycling symmetric and involves in the nearest neighbor sites instead of the next nearest neighbors.

Strong coupling limit. We consider H_T and H_{SSO} as the perturbations. Besides the well-known AFM coupling associated with H_T, the H_{SSO} contribution, to the second order for a given link a , is given by,

$$-\frac{2}{U}(H_{SSO})_{a,ij}(H_{SSO})_{a,ji} = -\frac{2}{U}(i\lambda\nu_{ij}c^\dagger_{i\sigma}\sigma^a_{\sigma\sigma'}c_{j\sigma'})(i\lambda\nu^a_{ji}c^\dagger_{j\tau}\sigma^a_{\tau\tau'}c_{i\tau'}) = \tilde{J}S^a_iS^a_j - \tilde{J}S^b_iS^b_j - \tilde{J}S^c_iS^c_j + \frac{\tilde{J}}{4}n_in_j - \frac{\tilde{J}}{2}n_i \quad (2)$$

where $\tilde{J} = \frac{4\lambda^2}{U}$, $a \neq b \neq c$ and $\mathbf{S} = \frac{1}{2}c^\dagger_\sigma\vec{\sigma}_{\sigma\sigma'}c_{\sigma'}$ is the electron spin operator. At the half filling, the SSO-Hubbard model reduces to

$$H_{J\tilde{J}} = J\sum_{\langle ij \rangle}\mathbf{S}_i\cdot\mathbf{S}_j + \tilde{J}\sum_{a;\langle ij \rangle_a}(S^a_iS^a_j - S^b_iS^b_j - S^c_iS^c_j), \quad (3)$$

where $J = \frac{4t^2}{U}$ as well-known. The crucial difference here from that in the KH model is both J and $\tilde{J}$ are positive, which means we have two different AFM orders. For $\lambda = 0$, it is AFM Heisenberg model. If $t = 0$, it is another quantum liquid which differs a sign from the stripy AFM model and is of a 'zig-zag' AFM order (See Fig. 1). It seems to be the most likely spin order in A_2IrO_3 in recent experimental reports combining with the first principles calculation [6–8] or a direct measurement [9]. This 'zig-zag' AFM model in fact can be mapped to an AFM model $H_{\tilde{J}} = \tilde{J}\sum_{\langle ij \rangle}\tilde{\mathbf{S}}_i\cdot\tilde{\mathbf{S}}_j$ with $\tilde{\mathbf{S}}_i$ defined as follows: Divide the honeycomb lattice into four sublattices, keep $\tilde{\mathbf{S}}_i = \mathbf{S}_i$ in one sublattice and change in other three, e.g., $\tilde{\mathbf{S}}_i = (S^x_i, -S^y_i, -S^z_i)$ for the x-link direction sublattice (and cycling for the y- and z-link's. See Fig. 1(a) [4].)



(a) Stripy AFM (b) Zig-zag AFM

FIG. 1: (Color online) (a) The stripy AFM order. Four different colored spots label four sublattices. (b) The zig-zag AFM order. The black and white spots divide the lattice in to a bipartite one.

Rewrite Eq. (3) as

$$H_{J\tilde{J}} = (J - \tilde{J})\sum_{\langle ij \rangle}\mathbf{S}_i\cdot\mathbf{S}_j + 2\tilde{J}\sum_{a;\langle ij \rangle_a}S^a_iS^a_j \quad (4)$$

$$= (\tilde{J} - J)\sum_{\langle ij \rangle}\tilde{\mathbf{S}}_i\cdot\tilde{\mathbf{S}}_j + 2J\sum_{a;\langle ij \rangle_a}\tilde{S}^a_i\tilde{S}^a_j. \quad (5)$$

One immediately sees that if $t = \lambda$, we arrives at Kitaev model with $J_K = 2J$. It is coaction both of the hopping and the SSO coupling gives birth to the Kitaev coupling. Notice that the system, either for $\tilde{J} = 0$ or $J = 0$, is in the conventional Landau symmetry breaking ordered phase. The Kitaev phase, however, is a SL phase without any Landau-type order parameter. That is to say, the Kitaev physics is the result of competition between two different AFM orders. There are quantum phase transitions from the Neel phase to the Kitaev SL and then to the 'zig-zag' AFM phase. The ground state energy per site for the Neel order is estimated as $e_N/J \approx (1 - \beta/3)\langle\mathbf{S}_i\cdot\mathbf{S}_j\rangle$ for $\beta = \tilde{J}/J$ and $\langle\mathbf{S}_i\cdot\mathbf{S}_j\rangle \approx -0.37$ [4] while the upper boundary of that of the Kitaev SL is $e_K/J \lesssim (1 + \beta)\langle S^a_iS^a_j\rangle$ for $\langle S^a_iS^a_j\rangle = -0.13$ [25]. This estimates the critical point $\beta_c \lesssim 0.95$. According Eq. (5), another critical point is $1/\beta_c \approx 1.05$. By exact diagonalization calculations, up to 24 lattice sites, we confirm the critical point is very close to the above estimate but are unable to determine that according to the ground state properties. Since the gapless excitation in the Kitaev phase is very different from that in the AFM phases: The former is Majorana fermion while the latter is bosonic spin wave. Therefore, we can estimate the critical points by looking for the level crossing of the first excitation state energy levels of two different phases in a finite size (See Fig. 2 (a)). (We also use this criteria to check the critical points in the KH model which are the same as the results in Ref. [4].) The 24-site data gives $\beta_c \approx 0.93$ for the phase transition from the Neel order to the Kitaev phase.

In Kitaev model, if a perturbed external magnetic field is applied, the SL becomes gapped and the phase we are discussed turns to be topologically non-trivial with a non-vanishing Chern number and non-abelian anyons. We expect these non-trivial properties exist in the whole SL phase and a gapless chiral excitation emerges in the edge of the system. Both of the AFM states are also gapped. At the critical points, the gaps collapse.

When the system deviates from the half filling, one can study the doped t-λ-J-$\tilde{J}$ model.

Original Kitaev model allows a phase transition from



FIG. 2: (Color online) (a) The level crossing of the energy levels of the first excited states for lattice sites $N = 8, 16$ and 24. (b) Schematic phase diagram in J-$\tilde{J}$ plane.

the non-abelian phase to three equal gapped SL phase with abelian anyon. This can also be realized in this SSO-Hubbard model by introducing link-dependent hopping amplitudes t^x, t^y, t^z and SO couplings $\lambda^x, \lambda^y, \lambda^z$. In this way, we can have a whole parameter Kitaev phase as well as the fruitful phase diagram from the XYZ model to Kitaev model and that in Kitaev model.

In the KH model studied in Ref. [4], there is a phase transition from the stripy AFM phase to the Kitaev phase. This can not be explained by the single band SSO repulsive Hubbard model. Instead, we find that it happens when two different FM orders [26] compete, i.e., if $J < 0$ and $\tilde{J} < 0$ in Eq. (4), one has

$$H'_{J\tilde{J}} = (|\tilde{J}| - |J|) \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j - 2|\tilde{J}| \sum_{a;\langle ij \rangle_a} S_i^a S_j^a. \quad (6)$$

(Or expressed by $\tilde{\mathbf{S}}$ with $\tilde{J} \leftrightarrow J$ as Eq. (5).) When $\tilde{J} = 0$, one has the FM order while it is the stripy AFM order [4] for $J = 0$. Again, Kitaev model comes out at $J = \tilde{J}$. The phase transition from the Kitaev phase to the stripy AFM phase happens at $J = 2\tilde{J}/3$ according to the result by Ref. [4]. Thus, the window of the Kitaev phase is larger than the AFM case: $2/3 \le \tilde{J}/J \le 3/2$. This captures the part of the KH model [4] from the stripy AFM to the Kitaev phases. Furthermore, the phase transition from the AFM Neel order to the stripy AFM order [4] is the result of the competition between $J > 0$ and $\tilde{J} < 0$. A model with $J < 0$ and $\tilde{J} > 0$ corresponds to a phase transition from the FM to the zig-zag AFM order. A schematic phase diagram is depicted in Fig. 2(b).

Weak interaction limit. When $U = 0$, the free-electron gas on honeycomb lattice has a gapless Dirac fermonic low energy spectrum in two valleys near the Dirac points. Now we discuss the effect of the SSO coupling to the Dirac semi-metal. The honeycomb lattice is bipartite to A and B sublattices (See Fig. 1(b)). Using the z-link as the elemental cell, the electron annihilation operator may be labeled by the sublattice suffices, i.e., $c_{\sigma i} \to (c_{A\sigma,i}, c_{B\sigma,i})$. Under the Fourier transformation $c_{\alpha\sigma} = \frac{1}{\sqrt{2N}} \sum_{\mathbf{k}} c_{\alpha\sigma\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}_i}$, the Hamiltonian with the SSO coupling is given by

$$H_T + H_{SSO} = \sum_{\mathbf{k}} (c^\dagger_{A\uparrow\mathbf{k}}, c^\dagger_{B\uparrow\mathbf{k}}, c^\dagger_{A\downarrow\mathbf{k}}, c^\dagger_{B\downarrow\mathbf{k}}) \begin{pmatrix} 0 & -tf(k) + i\lambda & 0 & \lambda g_-(k) \\ -tf^*(k) - i\lambda & 0 & \lambda g_+(k)^* & 0 \\ 0 & \lambda g_+(k) & 0 & -tf(k) - i\lambda \\ \lambda g_-(k)^* & 0 & -tf^*(k) + i\lambda & 0 \end{pmatrix} \begin{pmatrix} c_{A\uparrow\mathbf{k}} \\ c_{B\uparrow\mathbf{k}} \\ c_{A\downarrow\mathbf{k}} \\ c_{B\downarrow\mathbf{k}} \end{pmatrix} \quad (7)$$

where $f(k) = 1 + e^{i\mathbf{k}\cdot\mathbf{n}_1} + e^{i\mathbf{k}\cdot\mathbf{n}_2}$ and $g_\pm(k) = i(e^{i\mathbf{k}\cdot\mathbf{n}_1} \pm ie^{i\mathbf{k}\cdot\mathbf{n}_2})$ with the vectors $\mathbf{n}_1 = (1/2, \sqrt{3}/2)$ and $\mathbf{n}_2 = (-1/2, \sqrt{3}/2)$. The spectra of quasi-electron excitations then can be easily calculated:

$$E_\pm^2 = (6\lambda^2 + 2t^2|f|^2)/2 \quad (8)$$
$$\pm\sqrt{(6\lambda^2 + 2t^2|f|^2)^2 - 4|\lambda^2 g_+ g_- - t^2 f^2 - \lambda^2|^2}/2$$

When $\lambda = 0$, we recove $E = \pm t|f(k)| \propto q$ near Dirac points $\mathbf{K}_\pm = (\pm 4\pi/3, 0)$ ($\mathbf{q} = \mathbf{k} - \mathbf{K}_\pm$). Now, because $|f|^2 \propto q^2$ and $|g_+(k)g_-(k) - 1| \propto q$ near $\mathbf{K}_\pm$, we have

$$E_- \approx \pm\lambda|g_+ g_- - 1| \sim \pm\lambda q, \; E_+ = \pm\sqrt{6}\lambda + O(q^2). \quad (9)$$

The t-dependent branch (E_+) of fermnionic excitation is gapped while a new gapless Dirac fermion mode (E_-) with a speed λ appears at the same Dirac points. This Dirac fermion differs from the free one as the latter does not mix the spins and the former mixes the spins.

Moderate interaction. In a moderate U, it is predicted that there is a featureless gapped SL phase [19]. A slave boson study showed that there is also a charge-spin gapped phase in a moderate U region [22] although the magnitude of this region is numerically different that from the quantum Monte Carlo study in Ref. [19]. Moreover, the SL state is broken down due to translation symmetry and lattice rotation symmetry breaking from a staggered $U(1)$ fluctuation. As we have seen before new physical properties of the system arise from the SSO interaction. We now use the salve boson technique to see what is the effect of this SSO coupling to the system in this interaction region.

We briefly review the slave boson theory [20, 21]. There are four different states in a single site: $|0\rangle, |\uparrow\rangle, |\downarrow\rangle, |\uparrow\downarrow\rangle$ with the completeness $|0\rangle\langle 0| + |\uparrow\rangle\langle\uparrow| + |\downarrow\rangle\langle\downarrow| + |\uparrow\downarrow\rangle\langle\uparrow\downarrow| = 1$. The electron operator can be expressed as $c_\sigma = |0\rangle\langle\sigma| + \sigma|-\sigma\rangle\langle\uparrow\downarrow|$. At half-filling, one can thing each site has averagely an electron charge. If subtracting this average charge, the state $|0\rangle$ can be thought as carrying a positive charge while $|\uparrow\downarrow\rangle$ carries a negative charge and $|\sigma\rangle$ carries spin degree with charge neutral. In the operator formalism, one has $c_\sigma = h^\dagger f_\sigma + \sigma f^\dagger_{-\sigma} d$. This is the slave boson technique with a spin-charge separation if $h^\dagger \sim |0\rangle, d^\dagger \sim |\uparrow\downarrow\rangle$, called holon and doublon, are bosonic operators carrying charge ± while $f^\dagger_\sigma \sim |\sigma\rangle$ is called charge neutral fermonic spinon. The completeness condition becomes a constraint on these charge and spin degrees: $s_i = n_{hi} + n_{di} + \sum_\sigma n_{f\sigma i} - 1 = 0$ for every site. Within the slave boson theory, then, the effective Hamiltonian is given by

$$\begin{aligned} H_{eff} &= U\sum_i d_i^\dagger d_i - \sum_a \sum_{a-links} [(t\chi^0_{f,ij} - \lambda\chi^a_{f,ij})\chi_{b,ij} \\ &+ (t\Delta^{0\dagger}_{f,ij} - \lambda\Delta^{a\dagger}_{f,ij})\Delta_{b,ji} + h.c.] + i\sum_i \xi_i s_i, \quad (10) \end{aligned}$$

where $\chi^0_{f,ij} = \sum_\sigma f^\dagger_{i\sigma} f_{j\sigma}$, $\chi^a_{f,ij} = \sum_{\sigma\sigma'} \sigma'\nu_{ij} f^\dagger_{i\sigma}\sigma^a_{\sigma\sigma'} f_{j\sigma}$, $\chi_{b,ij} = h_i^\dagger h_j - d_i^\dagger d_j$. And $\Delta^0_{f,ij} = \sum_\sigma \sigma f_{i,-\sigma} f_{j\sigma}$, $\Delta^a_{f,ij} = \sum_{\sigma\sigma'} \sigma'\nu_{ij} f_{i\sigma}\sigma^a_{\sigma\sigma'} f_{j,-\sigma'}$ and $\Delta_{b,ij} = h_i d_j + d_i h_j$. For a non-zero Δ_f then Δ_b, the pair nature of the boson wave function and the holon-doublon symmetry require $\chi_b = 0$ and then $t\chi^0_f - \lambda\chi^a_f = 0$.

Now we follow the same trick used in Ref. [22] to solve the mean field theory. The boson problem in our model is not much different from that in Ref. [22] if $t\Delta^0_f$ in Ref. [22], the mean field value of $\Delta^0_{f,ij}$, is replaced by $t\Delta^0_f - \lambda\Delta^a_f$ in every link. The charge excitation spectrum is $E^\pm_b = \pm\frac{U}{2} + \sqrt{(\frac{U-2\xi_0}{2})^2 - |\Delta_{fk}|^2}$ where $\xi_0 = \langle i\xi\rangle$ and $\Delta_{fk} = \sum_a (t\Delta^0_f - \lambda\Delta^a_f) e^{i\mathbf{k}\cdot\mathbf{n}_a}$ for $\mathbf{n}^{x,y} = \mathbf{n}_{1,2}$ and $\mathbf{n}^z = 0$ [22]. The mean field effective Hamiltonian for spinon is given by

$$\begin{aligned} H_f &= -(\xi_0/2)\sum_k (f^\dagger_{A\sigma,k} f_{A\sigma,k} + f^\dagger_{B\sigma,k} f_{B\sigma,k}) \quad (11) \\ &+ \sum_k \Delta_b [f^\dagger_{A\uparrow,k} f^\dagger_{B\downarrow,-k}(-tf(k) + i\lambda) \\ &+ f^\dagger_{A\downarrow,k} f^\dagger_{B\uparrow,-k}(tf(k) + i\lambda)] + \sum_k [f^\dagger_{A\uparrow,k} f^\dagger_{B\uparrow,-k}\lambda g_-(k) \\ &- f^\dagger_{A\downarrow,k} f^\dagger_{B\downarrow,-k}\lambda g_+(k)]\Delta_b + h.c. \end{aligned}$$

The eigen energies then are easily obtained by

$$\begin{aligned} E^2 &= \xi_0^2/4 + [6\tilde\lambda^2 + 2\tilde t^2|f|^2 \quad (12) \\ &\pm(1/2)\sqrt{(6\tilde\lambda^2 + 2\tilde t^2|f|^2)^2 - 4|\tilde\lambda^2 g_+ g_- - \tilde t^2 f^2 - \tilde\lambda^2|^2}] \end{aligned}$$

where $\tilde t = t\Delta_b$ and $\tilde\lambda = \lambda\Delta_b$. Notice that in Eq. (12), except the first term $\xi_0^2/4$, the rest is of the same form as Eq. (8), that of the Dirac fermions at $U = 0$. Therefore, there are two branches of spinon excitations whose energy minima are at the Dirac points and the gaps are $|\xi_0|/2$ and $\sqrt{\xi_0^2/4 + 6\tilde\lambda^2}$, respectively. As the same as that in Ref. [22], our mean field states also suffer a staggered $U(1)$ global gauge fluctuation as $f_{iA} \to e^{i\alpha} f_{iA}$ and $f_{iB} \to e^{-i\alpha} f_{iB}$ and the SL state is possibly broken down in a semiclassical limit. Besides the next nearest neighbor hopping may destroy this U(1) gauge symmetry [22], the Kane-Mele SO coupling [24] also concerns the next nearest neighbor sites and breaks this U(1) to Z_2 gauge symmetry so that the system is turned to be a SL. Moreover, while the Kane-Mele SO coupling gives a mass to the new Dirac fermion at the weak coupling and the system is expected to be a topological insulator with the time reversal symmetry[24], the strongly-corrected behaviors with this SO coupling are more intriguing [27]. Finally, if the system is doped, those SL phases may be superconductors and the competition between the d-wave superconductivity and a p-wave superconductivity is expected as U varys.

In conclusions, we have studied the physical behaviors of the SSO-Hubbard model in the strong, weak and moderate repulsive interactions. We showed that a SL phase can be a result of a competition between two Landau-type ordered states: the Kitaev phase appears when two different AFM couplings become comparable, i.e., both the hopping amplitude and the SO coupling play equal roles. Meanwhile, the hopping does not play any role in the limit of $U \to 0$. A phase with both charge and spin gapped emerges in the midway between the weak and strong couplings. We do not determine the critical U for these different phases. Since our mean field equation for the slave boson's is formally the same as that given in Ref. [22], the result is the same. The critical interaction into the AFM orders is also very close that when $\lambda = 0$. We have seen that the Kitaev phase appears due to the competition between two different AFM orders. This implies the Kitaev phase must not appear before these two AFM phases but once these two ordered ones exist, there should be this SL as a result of the competition.

In conclusions, we have studied the SSO-Hubbard model for the repulsive Hubbard U. This work was supported by NNSF of China and the 973 program of MOST of China.

[1] A. Kitaev, Ann. Phys. **321**, 2(2006).

[2] A. Kitaev, Ann. Phys. **303**, 2 (2003).

[3] L-M. Duan, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **91**, 090402 (2003).

[4] J. Chaloupka, G. Jackeli, and G. Khaliullin, Phys. Rev. Lett. **105**, 027204 (2010).

[5] G. Jackeli, and G. Khaliullin, Phys. Rev. Lett. **102** 017205 (2009).

[6] X. Liu, T. Berlijn, W.-G. Yin, W. Ku, A. Tsvelik, Y-J. Kim, H. Gretarsson, Y. Singh, P. Gegenwart, and J. P. Hill1, Phys. Rev. B **83**, 220403(R) (2011).

[7] Y. Singh, S. Manni, J. Reuther, T. Berlijn, R. Thomale, W. Ku, S. Trebst, and P. Gegenwart, arXiv:1106.0429v2 (2011).

[8] S. K. Choi et al, arXiv: 1202.1268, to appear in Phys. Rev. Lett.

[9] F. Ye, S. X. Chi, H. B. Cao, B. C. Chakoumakos, J. A. Fernandez-Baca, R. Custelcean, T. F. Qi, O. B. Korneta, and G. Cao, arXiv:1202.3995.

[10] F. Trousselet, G. Khaliullin, and P. Horsch, Phys. Rev. B **84**, 054409 (2011).

[11] J. Reuther, T. Thomale, and S. Trebst, Phys. Rev. B **84** 100406 (2011).

[12] H.-C. Jiang, Z.-C. Gu, X.-L. Qi, and S. Trebst, Phys. Rev. B **83**, 245104 (2011).

[13] J. Kimichi and Y-Z. You, Phys. Rev. B **84**,180407(2011).

[14] W-L. You and Y-L. Dong, Phys. Rev. B **84**, 174426 (2011).

[15] S. Bhattacharjee, S-S. Lee, and Y. B. Kim, arXiv: 1108.1806 (2011).

[16] Y-Z. You, I. Kimchi, and A. Vishwanath, Arxiv 1109.4155 (2011) .

[17] T. Hyart, A. R. Wright, G. Khaliullin, and B. Rosenow, Arxiv 1109.6681 (2011).

[18] G. Khaliullin, Prog. Theor. Phys. Suppl. **16**, 155 (2005).

[19] Z. Y. Meng, T. C. Lang, S. Wessel, F. F. Assaad and A. Muramats, Nature **464**, 847 (2010).

[20] Z. Zou and P. W. Anderson, Phys. Rev. B **37**, 627 (1988).

[21] P. A. Lee, N. Nagaosa, and X-G. Wen, Rev. Mod. Phys. **78**, 17 (2006).

[22] A. Vaezi and X-G. Wen, arXiv: 1010. 5744 (2010).

[23] S-L. Yu, X. C. Xie, and J-X. Li, Phys. Rev. Lett. **107**, 010401 (2011).

[24] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 146802 (2005); **95**, 226801 (2005).

[25] G. Baskaran, S. Mandal, and R. Shankar, Phys. Rev. Lett. **98**, 247201 (2007).

[26] We do not investigate the origin of these two FM phases in this work.

[27] L. Liang, S. J. Qin, and Y. Yu, in preparation.